7/18/25, 4:12 PM District Advisory



# Gramin Krishi Mausam Sewa Agromet Advisory Bulletin for KANNUR District Kerala Agricultural University



Vellanikkara, Thrissur, Kerala

# **Agromet Advisory Bulletin**

Date: 2025-07-18

Weather Forecast of District KANNUR (Kerala) Issued On: 2025-07-18 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-07-19	2025-07-20	2025-07-21	2025-07-22	2025-07-23
Rainfall(mm)	40.0	35.0	40.0	25.0	25.0
Tmax(°C)	29.0	29.0	29.0	29.0	29.0
Tmin(°C)	23.0	23.0	23.0	23.0	23.0
RH-I(%)	95	95	95	95	95
RH-II(%)	90	90	90	90	90
Wind Speed(kmph)	5	4	3	3	3
Wind Direction(Degree)	320	290	290	290	290
Cloud Cover(Octa)	8	8	8	8	8
Warning	Extremely Heavy Rain; Strong Surface Winds	Extremely Heavy Rain; Strong Surface Winds	Extremely Heavy Rain; Strong Surface Winds	Very Heavy Rain; Strong Surface Winds	Heavy Rain; Strong Surface Winds

### **Forecast Summary:**

The sky will be cloudy. High relative humidity will be experienced. The atmospheric temperature will come down. There will be extremely heavy rainfall on July19&20. There will be heavy to very heavy rainfalls on July21 in isolated places. There will be heavy rainfalls on July22 in isolated places.

### Weather Warnings (Valid Till 08:30 IST of the next day)

Chances for extremely heavy rainfall on July18.

### Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Difficulty in drying agricultural produce in open conditions. Chances for fall off of slender stemmed fruits and vegetables. Chances for water stagnation in the fields.

#### **General Advisory:**

Drain the stagnating water from the cultivated areas where heavy water stagnations occurred due to the last five days heavy rainfalls Harvest all the ripened products if any quickly. Do not keep the harvested products in the field itself. Do not attempt to dry the products like rice, rubber, copra, pulses, vegetable seeds, etc directly under sun. Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc. Clean the drainage channels in crop lowlands to enable proper drainage of excess water in case of exigencies if any. Do not shelter the cattle in houses with weak structures. Keep animal feeds, seeds, fertilizers etc. in termite free and

7/18/25, 4:12 PM District Advisory

moist free rooms on elevated platforms coated with wooden panels. Avoid activities under open conditions while there is lightning. Also do not expose livestock to such unfavorable weather conditions. Seedlings of tree crops can be planted now. Ensure proper distance between the plants. The 2/3 portions of the pits should be filled with mixture of topsoil and powdered dry cow dung/compost. Spraying should be done only if sufficient intervals are available between the rainfalls. Use sticking agents like

### **SMS Advisory:**

Black pepper: Pollu beetle: Mix quinalphos 25%EC (@ 2ml per litre) with 1% of Bordeaux mixture and spray

### **Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory
RICE (Tillering)	Leaf folder: If leaf folder attack is seen, unfold the leaves by rolling thorny wires/twigs over the canopy. If the attack is severe spot spraying of flubendiamide @ 1.0 ml/10L is recommended. Attack will be more in shaded conditions. Avoid excessive Nitrogenous fertilizers

### **Horticulture Specific Advisory:**

Horticulture (Stage)	Horticulture Specific Advisory
COCONUT (Seedling Growth)	Water stagnation and decaying of growing bud (bud rot): Do not allow water stagnation in the planted pits. Provide bunds around the pits to prevent the collection of runoff water in the pits. After the cessation of heavy rainfall, spray 1 % Bordeaux Mixture on the leaves and growing buds, only after mixing with a sticking agent to the fungicide.
COCONUT (Grand growth)	Rhinoceros beetle: Take out the beetles from the attacked palm crown using beetle hook. Swab the hole made by the beetles with Bordeaux paste to prevent entry of fungus through the cut surface. Old fishnets can be reused for controlling the pest. Cut the nets into pieces of size $1 \text{m x} 0.5 \text{m}$ . Keep the middle portion of the net, lengthwise into the axils of 3rd and 4th leaf whirls of the palm. Push the net firmly with small stones into the axils. The hanging sides of the net should be kept open. The beetles will be trapped in the nets and get destroyed. Care should be taken to move the nets to upper leaf axils accordingly while fresh leaves are emerging.
BLACK PEPPER (Grand growth)	Pollu beetle: Mix quinalphos 25%EC (@ 2ml per litre) with 1% of Bordeaux mixture and spray
BLACK PEPPER (Grand growth)	Foot rot: As prophylactic measure, apply 150 gram of Trichoderma enriched neem cake - cow dung mixture in the basins of the vines and incorporate thoroughly with the soil. If disease already appeared, drench soil in the plant basins with Redomil 0.2% (2g/litre of water). Spray the same on the leaves also.
BANANA (Grand growth)	Rhizome rot: Ensure good drainage in the garden Drench with Streprocyclin antibiotic (@2g/10L) Or Copperhydroxide (20g/10L) at the basin of the plant.

### **Poultry Specific Advisory:**

Poultry	Poultry Specific Advisory	
CHICKEN	Cage hygiene: Keep the cages dry. If moisture persists, ammonia may be	
CHICKEN	generated from the droppings which will adversely affect the health of the birds.	

#### Likely impacts of weather warnings (General)

Visibility may become poor due to intense spell of rainfall leading to traffic congestion.
 Temporary
Disruption of traffic due to water logging in roads/ uprooting of trees/ breaking of tree branches leading to
increased travel time.
 Uprooting of trees may cause damages to power sector.
 Possibility of damages to

7/18/25, 4:12 PM District Advisory

vulnerable structures due to heavy to very heavy rainfall • Partial Damages to Kutcha Houses and Huts due to uprooting of trees. • Possibilities of Flash floods due to intense spell of rainfall. • Water logging / flooding in many parts of low-lying areas. • Landslides/mud slide/land slip very likely.

### Impact based advisories (General)

• Follow traffic advisories issued, if any. • Avoid staying in vulnerable structure. • Avoid going to the areas that face water logging problems/ river fronts. • People in the vulnerable area are advised to move to safer places.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam Mobile App link: https://play.google.com/store/apps/details/

Meghdoot Mobile App link: https://play.google.com/store/apps/details

Damini Mobile App link: https://play.google.com/store/apps/details