

Agromet Advisory Service Bulletin for Thiruvananthapuram District (Issued jointly by AMFU, RARS Vellayani, Kerala Agricultural University& India Meteorological Department) Based on Ensemble Weather forecast by IMD



Dated: 10.06.2025

Bulletin No. TVM-AAS-23/2025, (46)

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK

Rainfall(mm)	Max Temperature(	°C) Min Temper	Min Temperature(°C)		ative Humidity ('	%)	Evaporation(mm)		
10.5	32-32.2		22.4-23.5		70-91		3.5-4.5		
WEATHER FORECAST FOR THE NEXT FIVE DAYS									
Weather parameters		11/06/2025	12/06/2025		13/06/2025 14/		06/2025	15/06/2025	
Rainfall (mm)		8	10		12	15		20	
Maximum temperature(°C )		34	34		34	34		34	
Minimum temperature (°C )		26	26		26	26		26	
Maximum Relative Humidity (%)		85	85		85	85		85	
Minimum Relative Humidity (%)		75	75 75		75	75		75	
Wind speed (kmph)		10	10 6		3	8		12	
Wind direction (deg)		320	320		290	270		250	
Total cloud cover (octa)		8	8		8	8		8	
WEATHER SUMMERY ALERT		There is a chance of moderate rainfall in isolated places of the district. Cumulatively above normal rainfall is likely over kerala.							
GENERAL ADVISORY		Ensure proper drainage in the fields and provide propping for banana plants. Reinforce vegetable pandals adequately, and adopt proper mulching techniques to retain soil moisture							
		AGRO-METEORO	DLOGICAL	ADV	ISORIES				
		There is a cha	nce of she	ath	blight, sheath ro	ot, lea	af spot di	sease etc. As a	

Paddy (Virippu)	Land preparation to transplanting	There is a chance of sheath blight, sheath rot, leaf spot disease etc. As a prophylactic measure, either soak one kg of paddy seeds or dip root of seedlings in a solution of <i>Pseudomonas</i> culture @10gram of <i>Pseudomonas</i> culture per one litre of water for 30 minutes.			
Mango, Jackfruit, Cashew	Pink Disease	The disease appears like a spider web, followed by pink-colored powder on the branches. The affected branches dry up. To control this, apply 1% Bordeaux mixture on the affected areas.			
Coconut	Leaf blight	Yellowish brown spots with grey brown margins appear on leaflets causing extensive leaf blight. Remove the severely affected older leaves and burn it. Spray the crown with Bordeaux mixture 1% or Propiconazole 25 EC 2 ml/L.			
Livestock	Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospyrosis				