18/07/2025, 16:03 District Advisory



# Gramin Krishi Mausam Sewa Agromet Advisory Bulletin for THRISSUR District Kerala Agricultural University



Vellanikkara, Thrissur, Kerala

# **Agromet Advisory Bulletin**

Date: 2025-07-18

Weather Forecast of District THRISSUR (Kerala) Issued On: 2025-07-18 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-07-19	2025-07-20	2025-07-21	2025-07-22	2025-07-23
Rainfall(mm)	30.0	40.0	35.0	18.0	15.0
Tmax(°C)	28.0	28.0	28.0	28.0	28.0
Tmin(°C)	23.0	23.0	23.0	23.0	23.0
RH-I(%)	95	95	95	95	95
RH-II(%)	90	90	90	90	90
Wind Speed(kmph)	4	4	4	3	3
Wind Direction(Degree)	320	320	320	320	320
Cloud Cover(Octa)	8	8	8	8	8
Warning	Very Heavy Rain; Strong Surface Winds	Very Heavy Rain; Strong Surface Winds	Heavy Rain; Strong Surface Winds	Heavy Rain; Strong Surface Winds	Heavy Rain; Strong Surface Winds

#### **Forecast Summary:**

Weather summary / Alert: -. As per the forecast of IMD, there is a chance of moderate rainfall in the district and heavy rainfall in isolated places. Strong winds speed reaching likely to prevail occasionally over Kerala from 18th to 21st July 2025. Above normal rainfall is expected in Kerala.

#### Weather Warnings (Valid Till 08:30 IST of the next day)

Very Heavy Rain; Strong Surface Winds

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Strong winds speed reaching likely to prevail occasionally over Kerala from 18th to 21st July 2025

# **General Advisory:**

General Advisory: -Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. Pesticides should not be applied during rainy period. Provide supports for the bananas. Make the vegetable pandals stronger.

#### **SMS Advisory:**

As per the forecast of IMD, there is a chance of moderate rainfall in the district and heavy rainfall in isolated places.

### **Crop Specific Advisory:**

18/07/2025, 16:03 District Advisory

Crop (Stage)	Crop Specific Advisory		
RICE (Seedling Growth)	Rice (virippu)-Seedling stage- Fertilizer application- First dose fertilizers can be applied for short duration verities at the rate of 40 kg urea, 78 kg rock phosphate and 12 kg potash per acre.		
NUTMEG (Vagatativa)	Nutmeg -Leaf fall- During heavy rainfall proper drainage facilities have to be ensure and pesticides should not be applied. During rainy season, there is a chance of leaf fall in nutmeg, as a prophylactic measure spray 1% Bordeaux mixture or drench the basin.		

### **Horticulture Specific Advisory:**

Horticulture (Stage)	Horticulture Specific Advisory		
COCONUT (Vegetative)	Coconut -Management-Since sufficient soil moisture is available, coconut plants can be applied with apply 361 gram of urea, 592 gram of rock phosphate and 668 gram of muriate of potash per matured coconut plant.		
(Flowering)	Vegetables (Cowpea)- Anthracnose disease- During rainy season, there is a chance of Anthracnose disease in Cowpea. To control spray Carbendazim one gram per one litre of water. Pesticides should not be applied during rain.		

## **Live Stock Specific Advisory:**

Live Stock	Live Stock Specific Advisory		
COW	Animal Husbandry: Animals should not be allowed to go to water logged areas in order to avoid anthrax and foot lesions. Care should be taken to prevent rat breeding in the premises of cattle shed to prevent leptospirosis. (CAADECCS, KVASU, Mannuthy)		

## Likely impacts of weather warnings (General)

Harvest the crop as early as possible. Provide proper drainage in the cropped field as heavy rainfall is expected. Propping of the banana plants recommended. Additional support of the pandals recommended. Keep away from open fields to avoid lightning stroke to livestock and farmers in the afternoon.Livestock / Poultry / Fishery: Keep the animals inside the shed during heavy rainfall / hailstorm and provide them balanced feed.

### Impact based advisories (General)

Heavy rains may cause flooding in the lowlands. Heavy rainfall results in the flooding of the cropped field which damages the crop in the maturity stage. High winds cause lodging of the banana plants. Those vegetables in pandals will be damaged due to high winds.

Farmers are advised to download Unified "Mausam" and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightening.

Mausam MobileApp link: https://play.google.com/store/apps/details/

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link: https://play.google.com/store/apps/details