

## GraminKrishiMausamSewa(GKMS)

## Wayanad

## **Agromet Advisory Bulletin for Wayanad District**

90

4

250

90

4

250

90

3

270

90

3

250

90

3

290



AAB No.:57/ 2025-2026

88

3.1

 $\mathsf{SSW}$ 

88

2.2

 $\mathsf{SSW}$ 

3.1

SSW

84

2.3

SSW

84

3.7

 $\mathsf{SSW}$ 

90

2.3

SSW

	Observed w (Dated 11.						WeatherParameters	WeatherForecast (Valid for 19.07.25 to 23.07.25)				
11/07	12/07	13/07	14/07	15/07	16/07	17/07	Date	19/07	20/07	21/07	22/07	23/07
17.2	11.6	1.3	7.1	13.2	3.2	96.4	Rainfall(mm)	42	32	30	20	15
24.2	24.3	24.3	24.4	26.2	26.4	24	Temp max (°C)	28	28	28	28	28
17.1	17.1	17	17.3	16.9	15.9	17.3	Temp min( <sup>0</sup> C)	23	23	23	23	23
93	93	96	95	96	96	96	RH-I(%)	96	96	96	96	96
22	89	22	84	84	90	90	RH-II(%)	90	90	90	90	90

8 8 8 8 8 Weather summary / Alert: Heavy rainfall and windspeed are expected for the coming weeks. So farmers are advised to take appropriate precautionary measures.

WindSpeed(km/hr)

Cloud Cover(octa)

Winddirection

90

2.8

NNW

to take appropriate p	Jiecaationary meas					
NameofCrop/Stage	Pest/Disease	AgrometAdvisory				
Rice (Nursery stage)		Transplanting cane be done. Provide good drainage facilities in nursery				
Cardamom ( Maturation to harvest)	Clum rot/Azhukal/c apsule rot	Apply mulch and clean the drainage channels. Farmers are advised to keep vigilance against Shoot/panicle borer				
Coconut (Different stages)	Red palm weevil Rhinoceros Beetle Stem bleeding Leaf rot	Improvedrainage facility. Avoid the application of fertilizers and plant protection chemicals.				
Arecanut (Nut development)	Mahali	Take necessary phyto-sanitary measures and provide proper drainage facilities				
Pepper (berry development)	Quick Wilt	Improvedrainage facility. Avoid the application of fertilizers and plant protection chemicals.				
Banana (Different stages)	Bunchy top Rhizome weevil	Farmers are advised to provide propping to standing banana crop as there is chance of high wind speed in the coming dates. Clean the drainage channels. As there is chance of Pseudo stem weevil infestation farmers are advised to adopt strict phyto-sanitary measures.				
Ginger/Turmeric (Vegetative)	Ginger blast Shoot borer Rot and wilt diseases	Improve drainage facility by providing proper drainage channels				
Coffee (Berry development)	White Stem borer Leaf rust	Improve drainage . Fertilizer application shall be postponed				
Animal Husbandry	Caring	Avoiding moisture in places where cattle feeds are being stored reduce the chances of food poison. livestock shed and premises should be kept clean by the proper disposal of food waste for avoiding <i>Leptospirosis</i> .				
This Agro Advisory Bulletin (AAS) is Prepared by Kerala Agricultural University, RARS,Ambalavayal based on the medium range weather forecast of India Meteorological Department(IMD),NewDelhi						